

| States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Percentages, Annual Averages Based on 2013 and 2014 NSDUHs | | | | | | |
|---|--------------------------|---------------|---------------|---------------|---------------|---------------|
| Table | Age Group | Quintile 1 | Quintile 2 | Quintile 3 | Quintile 4 | Quintile 5 |
| Table 1. Illicit Drug Use in the Past Month | 12 to 17 | 6.31 - 7.51 | 7.52 - 8.37 | 8.38 - 9.41 | 9.42 - 10.39 | 10.40 - 14.58 |
| | 18 to 25 | 14.14 - 17.72 | 17.73 - 19.19 | 19.20 - 22.34 | 22.35 - 25.22 | 25.23 - 33.84 |
| | 26 or Older | 4.28 - 6.21 | 6.22 - 6.85 | 6.86 - 7.90 | 7.91 - 9.60 | 9.61 - 14.21 |
| | 18 or Older | 5.76 - 7.88 | 7.89 - 8.56 | 8.57 - 9.68 | 9.69 - 11.86 | 11.87 - 17.06 |
| | 12 or Older | 5.82 - 7.82 | 7.83 - 8.55 | 8.56 - 9.67 | 9.68 - 11.67 | 11.68 - 16.82 |
| Table 2. Marijuana Use in the Past Year | 12 to 17 | 9.90 - 11.19 | 11.20 - 12.37 | 12.38 - 14.00 | 14.01 - 15.65 | 15.66 - 20.81 |
| | 18 to 25 | 22.65 - 27.32 | 27.33 - 29.18 | 29.19 - 32.31 | 32.32 - 36.50 | 36.51 - 46.28 |
| | 26 or Older | 6.17 - 7.62 | 7.63 - 8.43 | 8.44 - 9.77 | 9.78 - 12.55 | 12.56 - 17.69 |
| | 18 or Older | 8.87 - 10.58 | 10.59 - 11.59 | 11.60 - 12.90 | 12.91 - 15.65 | 15.66 - 21.83 |
| | 12 or Older | 8.97 - 10.72 | 10.73 - 11.58 | 11.59 - 13.01 | 13.02 - 15.61 | 15.62 - 21.70 |
| Table 3. Marijuana Use in the Past Month | 12 to 17 | 4.98 - 5.60 | 5.61 - 6.22 | 6.23 - 7.65 | 7.66 - 8.74 | 8.75 - 12.56 |
| | 18 to 25 | 11.55 - 15.76 | 15.77 - 16.99 | 17.00 - 19.74 | 19.75 - 23.17 | 23.18 - 31.24 |
| | 26 or Older | 3.30 - 4.42 | 4.43 - 5.12 | 5.13 - 6.13 | 6.14 - 8.49 | 8.50 - 12.45 |
| | 18 or Older | 4.76 - 6.17 | 6.18 - 6.75 | 6.76 - 7.96 | 7.97 - 10.40 | 10.41 - 15.17 |
| | 12 or Older | 4.81 - 6.11 | 6.12 - 6.70 | 6.71 - 7.90 | 7.91 - 10.18 | 10.19 - 14.93 |
| Table 4. Perceptions of Great Risk from Smoking Marijuana Once a Month | 12 to 17 | 15.72 - 19.76 | 19.77 - 22.51 | 22.52 - 24.68 | 24.69 - 26.52 | 26.53 - 32.75 |
| | 18 to 25 | 5.97 - 9.53 | 9.54 - 12.60 | 12.61 - 13.67 | 13.68 - 15.62 | 15.63 - 22.76 |
| | 26 or Older | 16.48 - 23.67 | 23.68 - 28.30 | 28.31 - 29.63 | 29.64 - 32.58 | 32.59 - 41.70 |
| | 18 or Older | 15.04 - 21.32 | 21.33 - 25.91 | 25.92 - 27.36 | 27.37 - 30.13 | 30.14 - 38.74 |
| | 12 or Older | 15.43 - 21.12 | 21.13 - 25.71 | 25.72 - 26.96 | 26.97 - 29.95 | 29.96 - 37.70 |
| Table 5. First Use of Marijuana (Marijuana Incidence) | 12 to 17 | 3.45 - 4.68 | 4.69 - 5.28 | 5.29 - 6.06 | 6.07 - 6.58 | 6.59 - 9.13 |
| | 18 to 25 | 4.69 - 6.43 | 6.44 - 6.90 | 6.91 - 8.04 | 8.05 - 9.01 | 9.02 - 12.91 |
| | 26 or Older ¹ | -- | -- | -- | -- | -- |
| | 18 or Older | 0.93 - 1.08 | 1.09 - 1.16 | 1.17 - 1.35 | 1.36 - 1.55 | 1.56 - 2.43 |
| | 12 or Older | 1.42 - 1.62 | 1.63 - 1.78 | 1.79 - 1.99 | 2.00 - 2.34 | 2.35 - 3.16 |
| Table 6. Illicit Drug Use Other Than Marijuana in the Past Month | 12 to 17 | 2.44 - 2.78 | 2.79 - 2.97 | 2.98 - 3.31 | 3.32 - 3.69 | 3.70 - 4.85 |
| | 18 to 25 | 4.64 - 5.46 | 5.47 - 6.27 | 6.28 - 6.81 | 6.82 - 7.29 | 7.30 - 11.43 |
| | 26 or Older | 1.64 - 2.22 | 2.23 - 2.45 | 2.46 - 2.72 | 2.73 - 3.05 | 3.06 - 4.73 |
| | 18 or Older | 2.08 - 2.78 | 2.79 - 3.03 | 3.04 - 3.41 | 3.42 - 3.71 | 3.72 - 4.93 |
| | 12 or Older | 2.11 - 2.79 | 2.80 - 3.01 | 3.02 - 3.40 | 3.41 - 3.69 | 3.70 - 4.92 |

| States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Percentages, Annual Averages Based on 2013 and 2014 NSDUHs | | | | | | |
|---|-------------|---------------|---------------|---------------|---------------|---------------|
| Table | Age Group | Quintile 1 | Quintile 2 | Quintile 3 | Quintile 4 | Quintile 5 |
| Table 7. Cocaine Use in the Past Year | 12 to 17 | 0.33 - 0.41 | 0.42 - 0.47 | 0.48 - 0.51 | 0.52 - 0.71 | 0.72 - 1.01 |
| | 18 to 25 | 1.90 - 3.16 | 3.17 - 3.67 | 3.68 - 4.18 | 4.19 - 5.54 | 5.55 - 8.49 |
| | 26 or Older | 0.66 - 0.84 | 0.85 - 1.01 | 1.02 - 1.30 | 1.31 - 1.52 | 1.53 - 3.26 |
| | 18 or Older | 0.96 - 1.21 | 1.22 - 1.41 | 1.42 - 1.67 | 1.68 - 2.13 | 2.14 - 3.64 |
| | 12 or Older | 0.90 - 1.14 | 1.15 - 1.32 | 1.33 - 1.57 | 1.58 - 2.01 | 2.02 - 3.47 |
| Table 8. Nonmedical Use of Pain Relievers in the Past Year | 12 to 17 | 3.77 - 4.18 | 4.19 - 4.44 | 4.45 - 4.75 | 4.76 - 5.22 | 5.23 - 6.15 |
| | 18 to 25 | 7.02 - 7.46 | 7.47 - 8.03 | 8.04 - 8.56 | 8.57 - 9.24 | 9.25 - 10.98 |
| | 26 or Older | 2.48 - 2.82 | 2.83 - 3.04 | 3.05 - 3.44 | 3.45 - 3.70 | 3.71 - 4.11 |
| | 18 or Older | 3.08 - 3.52 | 3.53 - 3.85 | 3.86 - 4.17 | 4.18 - 4.39 | 4.40 - 4.91 |
| | 12 or Older | 3.16 - 3.56 | 3.57 - 3.90 | 3.91 - 4.27 | 4.28 - 4.46 | 4.47 - 4.95 |
| Table 9. Alcohol Use in the Past Month | 12 to 17 | 6.76 - 9.90 | 9.91 - 10.81 | 10.82 - 11.44 | 11.45 - 12.86 | 12.87 - 14.63 |
| | 18 to 25 | 36.61 - 55.22 | 55.23 - 59.13 | 59.14 - 62.92 | 62.93 - 65.30 | 65.31 - 72.88 |
| | 26 or Older | 35.21 - 50.88 | 50.89 - 55.87 | 55.88 - 59.52 | 59.53 - 62.51 | 62.52 - 70.87 |
| | 18 or Older | 35.48 - 51.71 | 51.72 - 56.18 | 56.19 - 60.21 | 60.22 - 62.86 | 62.87 - 71.23 |
| | 12 or Older | 31.90 - 47.27 | 47.28 - 51.80 | 51.81 - 55.55 | 55.56 - 58.39 | 58.40 - 68.03 |
| Table 10. Binge Alcohol Use in the Past Month | 12 to 17 | 4.90 - 5.55 | 5.56 - 5.99 | 6.00 - 6.38 | 6.39 - 6.81 | 6.82 - 8.01 |
| | 18 to 25 | 26.08 - 34.42 | 34.43 - 37.37 | 37.38 - 41.27 | 41.28 - 43.07 | 43.08 - 50.12 |
| | 26 or Older | 15.45 - 20.61 | 20.62 - 22.02 | 22.03 - 23.28 | 23.29 - 24.19 | 24.20 - 32.78 |
| | 18 or Older | 17.44 - 22.35 | 22.36 - 24.30 | 24.31 - 25.86 | 25.87 - 26.87 | 26.88 - 35.60 |
| | 12 or Older | 15.91 - 20.87 | 20.88 - 22.74 | 22.75 - 24.07 | 24.08 - 25.09 | 25.10 - 33.98 |
| Table 11. Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week | 12 to 17 | 31.17 - 35.44 | 35.45 - 37.05 | 37.06 - 38.38 | 38.39 - 40.53 | 40.54 - 46.68 |
| | 18 to 25 | 22.20 - 27.60 | 27.61 - 29.07 | 29.08 - 32.40 | 32.41 - 36.10 | 36.11 - 41.96 |
| | 26 or Older | 35.20 - 37.94 | 37.95 - 40.19 | 40.20 - 42.18 | 42.19 - 44.04 | 44.05 - 48.42 |
| | 18 or Older | 34.15 - 36.58 | 36.59 - 38.52 | 38.53 - 40.80 | 40.81 - 43.05 | 43.06 - 47.42 |
| | 12 or Older | 34.23 - 36.50 | 36.51 - 38.33 | 38.34 - 40.57 | 40.58 - 42.75 | 42.76 - 46.90 |
| Table 12a. Underage Alcohol Use in the Past Month | 12 to 20 | 13.38 - 20.13 | 20.14 - 22.08 | 22.09 - 23.55 | 23.56 - 26.18 | 26.19 - 31.07 |
| Table 12b. Underage Binge Alcohol Use in the Past Month | 12 to 20 | 10.49 - 12.62 | 12.63 - 13.98 | 13.99 - 14.61 | 14.62 - 16.24 | 16.25 - 20.28 |

| States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Percentages, Annual Averages Based on 2013 and 2014 NSDUHS | | | | | | |
|---|-------------|---------------|---------------|---------------|---------------|---------------|
| Table | Age Group | Quintile 1 | Quintile 2 | Quintile 3 | Quintile 4 | Quintile 5 |
| Table 13. Tobacco Product Use in the Past Month | 12 to 17 | 5.00 - 6.66 | 6.67 - 7.70 | 7.71 - 8.99 | 9.00 - 9.86 | 9.87 - 12.46 |
| | 18 to 25 | 23.60 - 34.02 | 34.03 - 36.80 | 36.81 - 40.79 | 40.80 - 42.46 | 42.47 - 50.63 |
| | 26 or Older | 18.40 - 23.98 | 23.99 - 25.92 | 25.93 - 28.01 | 28.02 - 30.05 | 30.06 - 37.71 |
| | 18 or Older | 19.45 - 25.78 | 25.79 - 27.46 | 27.47 - 29.59 | 29.60 - 31.94 | 31.95 - 39.42 |
| | 12 or Older | 17.67 - 23.72 | 23.73 - 25.73 | 25.74 - 27.70 | 27.71 - 29.75 | 29.76 - 37.20 |
| Table 14. Cigarette Use in the Past Month | 12 to 17 | 3.66 - 4.60 | 4.61 - 5.54 | 5.55 - 6.26 | 6.27 - 7.03 | 7.04 - 8.64 |
| | 18 to 25 | 20.80 - 27.52 | 27.53 - 30.40 | 30.41 - 32.45 | 32.46 - 34.11 | 34.12 - 40.36 |
| | 26 or Older | 15.14 - 19.27 | 19.28 - 21.66 | 21.67 - 22.95 | 22.96 - 25.23 | 25.24 - 31.39 |
| | 18 or Older | 16.51 - 21.01 | 21.02 - 22.49 | 22.50 - 24.33 | 24.34 - 26.20 | 26.21 - 32.58 |
| | 12 or Older | 15.27 - 19.72 | 19.73 - 21.23 | 21.24 - 22.62 | 22.63 - 24.58 | 24.59 - 30.57 |
| Table 15. Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day | 12 to 17 | 57.60 - 61.85 | 61.86 - 63.52 | 63.53 - 64.91 | 64.92 - 66.67 | 66.68 - 71.65 |
| | 18 to 25 | 57.05 - 61.88 | 61.89 - 63.48 | 63.49 - 65.20 | 65.21 - 68.34 | 68.35 - 72.60 |
| | 26 or Older | 63.58 - 67.22 | 67.23 - 70.21 | 70.22 - 71.96 | 71.97 - 74.51 | 74.52 - 78.16 |
| | 18 or Older | 62.65 - 66.66 | 66.67 - 69.35 | 69.36 - 71.27 | 71.28 - 73.59 | 73.60 - 77.08 |
| | 12 or Older | 62.49 - 66.24 | 66.25 - 68.83 | 68.84 - 70.24 | 70.25 - 72.75 | 72.76 - 76.08 |
| Table 16. Alcohol Dependence or Abuse in the Past Year | 12 to 17 | 2.12 - 2.49 | 2.50 - 2.66 | 2.67 - 2.82 | 2.83 - 3.35 | 3.36 - 3.86 |
| | 18 to 25 | 10.13 - 11.60 | 11.61 - 12.66 | 12.67 - 13.39 | 13.40 - 14.37 | 14.38 - 17.17 |
| | 26 or Older | 4.61 - 5.46 | 5.47 - 5.78 | 5.79 - 6.13 | 6.14 - 6.59 | 6.60 - 9.19 |
| | 18 or Older | 5.51 - 6.40 | 6.41 - 6.82 | 6.83 - 7.10 | 7.11 - 7.93 | 7.94 - 10.20 |
| | 12 or Older | 5.23 - 6.03 | 6.04 - 6.37 | 6.38 - 6.69 | 6.70 - 7.50 | 7.51 - 9.78 |
| Table 17. Alcohol Dependence in the Past Year | 12 to 17 | 0.77 - 0.90 | 0.91 - 0.96 | 0.97 - 1.06 | 1.07 - 1.14 | 1.15 - 1.35 |
| | 18 to 25 | 4.57 - 5.03 | 5.04 - 5.33 | 5.34 - 5.88 | 5.89 - 6.66 | 6.67 - 7.12 |
| | 26 or Older | 2.33 - 2.61 | 2.62 - 2.81 | 2.82 - 2.99 | 3.00 - 3.14 | 3.15 - 4.15 |
| | 18 or Older | 2.67 - 2.97 | 2.98 - 3.18 | 3.19 - 3.44 | 3.45 - 3.67 | 3.68 - 4.55 |
| | 12 or Older | 2.51 - 2.80 | 2.81 - 2.96 | 2.97 - 3.21 | 3.22 - 3.41 | 3.42 - 4.35 |
| Table 18. Illicit Drug Dependence or Abuse in the Past Year | 12 to 17 | 2.84 - 3.01 | 3.02 - 3.27 | 3.28 - 3.52 | 3.53 - 3.86 | 3.87 - 4.50 |
| | 18 to 25 | 5.42 - 6.13 | 6.14 - 6.63 | 6.64 - 7.19 | 7.20 - 7.83 | 7.84 - 9.47 |
| | 26 or Older | 1.34 - 1.57 | 1.58 - 1.70 | 1.71 - 1.90 | 1.91 - 2.06 | 2.07 - 2.58 |
| | 18 or Older | 2.02 - 2.26 | 2.27 - 2.49 | 2.50 - 2.66 | 2.67 - 2.86 | 2.87 - 3.50 |
| | 12 or Older | 2.14 - 2.35 | 2.36 - 2.56 | 2.57 - 2.75 | 2.76 - 2.94 | 2.95 - 3.53 |

| States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Percentages, Annual Averages Based on 2013 and 2014 NSDUHs | | | | | | |
|---|-------------|---------------|---------------|---------------|---------------|---------------|
| Table | Age Group | Quintile 1 | Quintile 2 | Quintile 3 | Quintile 4 | Quintile 5 |
| Table 19. Illicit Drug Dependence in the Past Year | 12 to 17 | 1.54 - 1.70 | 1.71 - 1.80 | 1.81 - 1.89 | 1.90 - 2.01 | 2.02 - 2.30 |
| | 18 to 25 | 3.70 - 4.17 | 4.18 - 4.63 | 4.64 - 5.21 | 5.22 - 5.69 | 5.70 - 7.97 |
| | 26 or Older | 1.01 - 1.15 | 1.16 - 1.30 | 1.31 - 1.37 | 1.38 - 1.50 | 1.51 - 2.37 |
| | 18 or Older | 1.47 - 1.65 | 1.66 - 1.78 | 1.79 - 1.91 | 1.92 - 2.06 | 2.07 - 2.90 |
| | 12 or Older | 1.50 - 1.68 | 1.69 - 1.77 | 1.78 - 1.92 | 1.93 - 2.06 | 2.07 - 2.86 |
| Table 20. Dependence or Abuse of Illicit Drugs or Alcohol in the Past Year | 12 to 17 | 4.06 - 4.70 | 4.71 - 4.93 | 4.94 - 5.25 | 5.26 - 5.63 | 5.64 - 6.44 |
| | 18 to 25 | 14.09 - 14.81 | 14.82 - 16.73 | 16.74 - 17.69 | 17.70 - 19.17 | 19.18 - 21.15 |
| | 26 or Older | 5.67 - 6.63 | 6.64 - 6.95 | 6.96 - 7.37 | 7.38 - 7.87 | 7.88 - 11.01 |
| | 18 or Older | 7.18 - 7.99 | 8.00 - 8.44 | 8.45 - 8.86 | 8.87 - 9.38 | 9.39 - 12.51 |
| | 12 or Older | 6.95 - 7.68 | 7.69 - 8.16 | 8.17 - 8.52 | 8.53 - 9.12 | 9.13 - 12.14 |
| Table 21. Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year | 12 to 17 | 2.32 - 2.82 | 2.83 - 3.02 | 3.03 - 3.25 | 3.26 - 3.52 | 3.53 - 4.13 |
| | 18 to 25 | 5.29 - 5.76 | 5.77 - 6.24 | 6.25 - 6.54 | 6.55 - 7.23 | 7.24 - 8.08 |
| | 26 or Older | 1.18 - 1.36 | 1.37 - 1.48 | 1.49 - 1.60 | 1.61 - 1.75 | 1.76 - 2.29 |
| | 18 or Older | 1.83 - 2.03 | 2.04 - 2.18 | 2.19 - 2.33 | 2.34 - 2.47 | 2.48 - 3.21 |
| | 12 or Older | 1.91 - 2.13 | 2.14 - 2.25 | 2.26 - 2.42 | 2.43 - 2.55 | 2.56 - 3.25 |
| Table 22. Needing But Not Receiving Treatment for Alcohol Use in the Past Year | 12 to 17 | 1.96 - 2.36 | 2.37 - 2.50 | 2.51 - 2.68 | 2.69 - 3.05 | 3.06 - 3.53 |
| | 18 to 25 | 9.74 - 11.26 | 11.27 - 12.34 | 12.35 - 12.89 | 12.90 - 14.01 | 14.02 - 16.58 |
| | 26 or Older | 4.60 - 5.26 | 5.27 - 5.48 | 5.49 - 5.81 | 5.82 - 6.11 | 6.12 - 8.62 |
| | 18 or Older | 5.42 - 6.11 | 6.12 - 6.42 | 6.43 - 6.78 | 6.79 - 7.33 | 7.34 - 9.65 |
| | 12 or Older | 5.15 - 5.83 | 5.84 - 6.08 | 6.09 - 6.38 | 6.39 - 6.94 | 6.95 - 9.25 |
| Table 23. Serious Mental Illness in the Past Year | 18 to 25 | 3.71 - 4.26 | 4.27 - 4.45 | 4.46 - 4.84 | 4.85 - 5.10 | 5.11 - 6.43 |
| | 26 or Older | 3.06 - 3.75 | 3.76 - 4.09 | 4.10 - 4.43 | 4.44 - 4.76 | 4.77 - 5.56 |
| | 18 or Older | 3.25 - 3.78 | 3.79 - 4.17 | 4.18 - 4.50 | 4.51 - 4.85 | 4.86 - 5.50 |
| Table 24. Any Mental Illness in the Past Year | 18 to 25 | 16.29 - 18.39 | 18.40 - 19.72 | 19.73 - 20.61 | 20.62 - 21.37 | 21.38 - 26.49 |
| | 26 or Older | 15.35 - 16.82 | 16.83 - 18.17 | 18.18 - 19.39 | 19.4 - 20.00 | 20.01 - 22.34 |
| | 18 or Older | 16.03 - 17.04 | 17.05 - 18.44 | 18.45 - 19.33 | 19.34 - 20.25 | 20.26 - 22.66 |
| Table 25. Had Serious Thoughts of Suicide in the Past Year | 18 to 25 | 6.15 - 6.98 | 6.99 - 7.33 | 7.34 - 7.56 | 7.57 - 8.05 | 8.06 - 10.29 |
| | 26 or Older | 2.78 - 3.16 | 3.17 - 3.37 | 3.38 - 3.57 | 3.58 - 3.75 | 3.76 - 4.23 |
| | 18 or Older | 3.34 - 3.83 | 3.84 - 3.97 | 3.98 - 4.18 | 4.19 - 4.37 | 4.38 - 4.85 |

| States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Percentages, Annual Averages Based on 2013 and 2014 NSDUHs | | | | | | |
|---|-------------|-------------|--------------|---------------|---------------|---------------|
| Table | Age Group | Quintile 1 | Quintile 2 | Quintile 3 | Quintile 4 | Quintile 5 |
| Table 26. Major Depressive Episode in the Past Year | 12 to 17 | 8.69 - 9.92 | 9.93 - 10.46 | 10.47 - 11.11 | 11.12 - 11.95 | 11.96 - 14.62 |
| | 18 to 25 | 6.90 - 8.20 | 8.21 - 8.80 | 8.81 - 9.40 | 9.41 - 10.07 | 10.08 - 11.80 |
| | 26 or Older | 5.01 - 5.82 | 5.83 - 6.14 | 6.15 - 6.63 | 6.64 - 6.95 | 6.96 - 7.85 |
| | 18 or Older | 5.29 - 6.22 | 6.23 - 6.61 | 6.62 - 6.97 | 6.98 - 7.32 | 7.33 - 8.26 |
| -- Not available. | | | | | | |
| NOTE: This table shows the boundaries used to create the maps shown in the document named "2013-2014 NSDUH National Maps of Prevalence Estimates, by State" at http://www.samhsa.gov/data/ . Maps for the 18 or older age group are available only for the following measures: serious mental illness (SMI), any mental illness (AMI), serious thoughts of suicide, and depression (i.e., major depressive episode or MDE). | | | | | | |
| NOTE: State estimates (which are based on a survey-weighted hierarchical Bayes estimation approach) fall into one of five quintiles according to their ranking. Because 51 states were ranked for each measure, the middle quintile was assigned to 11 states, and the remaining quintiles were assigned 10 states each. In some cases, a "quintile" could have more or fewer states than desired because two (or more) states had the same estimate (to two decimal places). When such ties occurred at the "boundary" between two quintiles, all of the states with the same estimate were conservatively assigned to the lower quintile. Those states with the highest estimates for a given measure are in red, with the exception of the perceptions of risk measures, for which the lowest perceptions of great risk are in red. Those states with the lowest estimates are in white, with the exception of the perceptions of risk measures, for which the highest perceptions of great risk are in white. The upper and lower limits of each quintile shown in the map legend collectively define a continuum and are not necessarily the actual values of a particular state. Tables containing specific estimates for these states are available on the 2013-2014 NSDUH state web page at http://www.samhsa.gov/data/ . | | | | | | |
| ¹ The average annual incidence estimates for first use of marijuana were low for the 26 or older age group and had an abbreviated range; therefore, no state grouping was made for this age group. | | | | | | |
| Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013 and 2014. | | | | | | |